

PATIENT NUMBER:

MULTI-DAY ABORTION PATIENT INSTRUCTIONS

CARE COLORADO

**AT CARE COLORADO, WE STRIVE ON PROVIDING PATIENT-BASED CARE. OUR PHONE LINE IS
AVAILABLE 24 HOURS A DAY FOR ANY QUESTIONS OR CONCERNs. PLEASE DO NOT HESITATE TO
REACH OUT TO OUR OFFICE. THANK YOU FOR TRUSTING US WITH YOUR CARE AND WE WISH YOU
THE BEST FROM YOUR CARE TEAM!**

**1930 E ORMAN AVENUE
PUEBLO, CO 81004**

**PHONE: (719) 884-4070
TEXT: (719) 888-7118
FAX: (719) 888-7721**

HELLO AND THANK YOU



During the evening and next day, your cervix will begin to gradually open from the insertion of the Laminaria that will be placed this morning. Laminaria is a "seaweed" stick that absorbs your body fluid and expands. These will stay in overnight, please do not remove anything that is placed by our office. We also will place a gauze soaked in betadine solution, to keep everything clean from bacteria.

BETADINE SOLUTION MAY CAUSE MULTI-COLOR DRAINAGE-THIS IS NORMAL.

YOU WILL NEED TO WEAR A PAD WHILE THE LAMINARIA IS IN PLACE AT ALL TIMES

When the laminaria is placed overnight, some patients may have hard menstrual-like cramping, while others may have no cramping at all. During this process, nausea and vomiting may occur. You will be given medications to take around the clock for the cramping and nausea.

It is very important that you stick to the medication schedule to keep your discomfort under control.

The medications that we use all work together, so it is important to take them as instructed.

While you have the laminaria in place, you need to be active. ***The more you walk, the better the pressure will subside and will expand and dilate your cervix.*** We recommend that you stay active all day and go to bed when you normally would.

Following is the list of medications you will be taking home and a schedule of when to take them:

AT BEDTIME FOR PAIN:

GABAPENTIN 300MG-1 TABLET

NAUSEA:

ONDANESTRON 4MG
TAKE EVERY 4-6 HOURS
SMALL WHITE TABLET

PAIN:

ACETAMINOPHEN 325MG
TAKE EVERY 4-6 HOURS
1-2 TABLETS AS NEEDED

CRAMPING:

IBUPROFEN 800MG
TAKE VERY 8 HOURS
LARGE WHITE TABLET

FIRST DOSE:

FIRST DOSE:

FIRST DOSE:

START DAY

OVERNIGHT INSTRUCTIONS

IF AT ANY TIME YOU FEEL THERE IS A PROBLEM, YOU SHOULD CALL THE CLINIC IMMEDIATELY. THERE WILL BE SOMEONE AVAILABLE 24 HOURS A DAY TO ANSWER CALLS. IF FOR SOME REASON YOU CANNOT GET IN TOUCH WITH THE PERSON ON CALL, YOU SHOULD GO TO THE EMERGENCY ROOM

IF YOU CANNOT GET IN TOUCH WITH CARE, GO TO THE NEAREST EMERGENCY ROOM AND TELL THE ER PHYSICIAN THAT YOU ARE IN THE PROCESS OF HAVING AN ABORTION. ASK THE ER PHYSICIAN TO CONTACT DR. LEAH TORRES FROM CARE COLORADO AT:

(719)884-4070

1 ACTIVITIES

No strenuous activities should be done while the laminara is in place, as it may move the position of the laminaria. You will need to have a driver/escort to accompany you for the next day as you will be receiving IV sedation.

2 DIET

You may eat whatever you like until 4 hours before your procedure. At that time, you may have something light, not heavy or greasy.

3 ATTIRE

On the day of your procedure, please wear baggy, loose fitting clothing so it is easier for us to dress you after your procedure. It is also recommended that you wear underwear conducive for wearing a menstrual pad.

4 BREAST CARE

You should wear a tight bra/sports bra for 2 weeks. Wear the bra 24 hours a day. Only take it off to shower. While in the shower, have your back to the water. Although warm water running over your breasts may feel good, this may cause you to lactate (make milk). Do not rub or stimulate the breasts. Your breasts may become large and engorged. This can be quite painful. You may take Tylenol and use ice packs to relieve the pain.

AFTER-CARE



These instructions include how to reach us, after care instructions, activities to avoid, and general information that is important to read over and understand.

If you have any questions or concerns, please review this information pack and call at any time if needed at: **719-884-4070**.

PLEASE CALL IF YOU HAVE ANY OF THE FOLLOWING SYMPTOMS/ISSUES:

- You start bleeding so heavily that you soak through two thick sanitary pads every hour for 2 hours consecutively (4 large pads in 2 hours)
- You begin to have severe abdominal cramping not relieved by any medications provided or the tips listed below
- Temperature greater than 100.4 that persists after taking medication and you have checked in twice within the hour

If you are unable to get in contact with a CARE Colorado staff member or feel your health is in danger, go to the nearest Emergency Room immediately. If you feel comfortable doing so, tell the ER physician that you had an abortion and ask the ER physician to contact Dr. Leah Torres from CARE.

TAKE THE FOLLOWING MEDICATIONS THAT WERE PROVIDED TO YOU AS NEEDED

- **ACETAMINOPHEN 325MG - 2 TABLETS** (REPEAT EVERY 4-6 HOURS AS NEEDED FOR PAIN)
- **IBUPROFEN 200MG - 4 TABLETS** (REPEAT EVERY 8 HOURS AS NEEDED FOR PAIN)

FINISH DAY

WHAT TO EXPECT

BLEEDING/CLOTTING

BLEEDING AND CLOTTING ARE NORMAL, IT USUALLY OCCURS "OFF AND ON" AND COULD LAST FOR 4-12 WEEKS. IF YOU HAVE HAD A PRIOR C-SECTION(S) DELIVERY, YOU MAY EXPERIENCE HEAVIER BLEEDING.

MENSTRUAL PERIOD

YOUR NEXT PERIOD CAN BE 3-12 WEEKS AFTER YOUR ABORTION AND MAY BE HEAVIER THAN NORMAL. TAKING A FORM OF HORMONAL BIRTH CONTROL CAN HELP REGULATE YOUR CYCLE AND MAY HELP YOUR PERIOD RETURN QUICKER. SOME PEOPLE CAN OVULATE AS SOON AS 3 WEEKS AFTER A PREGNANCY ENDS SO USE PRECAUTIONS PRIOR TO YOUR PERIOD COMING BACK.

FEVER/TEMPERATURE ELEVATION

AFTER COMPLETION OF MISOPROSTOL, YOU CAN RUN A SLIGHT FEVER. IF YOUR FEVER IS 100.4 OR HIGHER FOR MORE THAN 8 HOURS, PLEASE CONTACT OUR OFFICE FOR FURTHER INSTRUCTIONS.

CRAMPING

CRAMPING IS COMMON AND MAY OCCUR "OFF AND ON" FOR 2-3 WEEKS AFTER COMPLETION OF ABORTION. TO REDUCE SYMPTOMS, TAKE ACETAMINOPHEN & IBUPROFEN (PROVIDED BY CLINIC) AS NEEDED. UTERINE MASSAGES ARE ALSO VERY IMPORTANT TO RECOVERY. THEY HELP WITH CRAMPING AND REDUCE BLEEDING RISK. HEATING PADS AND PHYSICAL ACTIVITY FOR 10-15 MINUTES WILL HELP REDUCE CRAMPS.

PREGNANCY SYMPTOMS

SIGNS OF PREGNANCY USUALLY DISAPPEAR WITHIN 48 HOURS. WHILE SYMPTOMS MAY HAVE GONE AWAY, IT IS STILL IMPORTANT COMPLETE FOLLOW-UP TO ENSURE THAT EVERYTHING IS COMPLETE.

ACTIVITIES TO AVOID

THERE ARE NO PHYSICAL RESTRICTIONS AFTER AN ABORTION. YOU CAN RETURN TO NORMAL ACTIVITIES THE DAY AFTER YOUR ABORTION.

QUESTIONS & CONCERNS



NEED TO KNOW

YOUR ABORTION IS YOUR BUSINESS AND WE WILL NOT DISCLOSE ANY OF YOUR INFORMATION

HAVING AN ABORTION DOES NOT AFFECT YOUR FUTURE FERTILITY

YOU DO NOT NEED TO TELL ANY HEALTH CARE PROVIDER YOU HAD AN ABORTION

In the event you experience any of the following, please see your nearest emergency room:

COMPLICATIONS:

HEAVY BLEEDING:

SOAKING 4 OVERNITE PADS FOR 2 HOURS CONSECUTIVELY, ALONG WITH FEELING FAINT

SEVERE ABDOMINAL CRAMPING:

SEVERE CRAMPING THAT CANNOT BE RELIEVED BY ANY MEDICATION PROVIDED

FEVER:

TEMPERATURE GREATER THAN 100.4 THAT PERSISTS AFTER TAKING MEDICATIONS

HOW MUCH AM I BLEEDING?




Please review this bleeding chart when calling about bleeding concerns. This helps us understand if this is a normal amount of bleeding or not, and can advise you further.



SCANT AMOUNT

BLOOD ONLY ON TISSUE
WHEN WIPED OR LESS
THAN ONE-INCH STAIN ON
MAXI PAD WITHIN ONE
HOUR



LIGHT AMOUNT

LESS THAN FOUR-
INCH STAIN ON MAXI
PAD WITHIN ONE
HOUR



MODERATE AMOUNT

LESS THAN SIX-
INCH STAIN ON
MAXI PAD WITHIN
ONE HOUR



HEAVY AMOUNT

SATURATED MAXI PAD
WITHIN ONE HOUR